



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

first found at Oxford has disappeared; so that the only station now known is that about two and a half miles south of the village of Oxford. The plants grow in shaded alluvial soil, on the bank of the Chenango River.

The Mode of Destruction of the Potato by *Peronospora infestans*.*

The tuber of the potato being only an enormous development of the stem must possess all the parts that characterize the exogenous stem. That is, it should show the central pith, the woody or vascular layer, the bark and the epidermis. These may be readily seen with the naked eye on slicing the potato. The vascular layer appears as a semi-transparent line running along the cut surface about a quarter of an inch below the cuticle and rising to the eyes where it meets the layer that represents the bark. At these points the cuticle is exceedingly thin and here the assault of *P. infestans* is usually made. Its progress is marked by a black streak advancing from the eye along the layer of vascular tissue which it destroys, either rising to the other eyes in its course or meeting similar hosts of enemies advancing from them. A layer of the bark immediately under the peel is attacked in the same manner about the same time. Two parallel black streaks are therefore visible advancing side by side below the skin of the potato.

In these two layers the life of the potato resides and their destruction consequently ensures its death. To them the attack is confined. The starch-laden cells of the pith and those between the vascular layer and the epidermis are not injured. Indeed a tuber in this or even in later stages of decay, is one of the best sources from which can be obtained perfect starch-cells with their contents uninjured for microscopic study.

Here the action of *P. infestans* ends. But other enemies soon troop in and complete the ruin it has begun. *Fusisporium Solani* abounds, but the greater part of what remains, the starch cells, is destroyed by *Bacterium termo*, the ubiquitous and ever-ready agent of destruction. This soon reduces the tuber to a mass of decay and to its action is principally due the smell arising from rotten potatoes. E. W. CLAYPOLE, AKRON, Ohio.

* Read before the Botanical Club of the A. A. A. S. at the Buffalo Meeting, August, 1886.